

SIDS

Notes by Craig Paardekooper

Looking at the **Eudra-vigilance** data, that covers 1504 different drugs, I applied the criteria for a safety signal, namely PRR > 2, with a lower confidence limit also greater than 2, and a minimum number of cases >= 3. The results are shown below. You can access this dataset here - <https://howbad.info/index.html#eudata>

| DRUG | SIDS | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|------------|------|------------|--------|-----------|-----------|---------|--------|
| BEYFORTUS | 5 | 523 | 0.956 | 14987141 | 240 | 0 | 597 |
| ROTASHIELD | 3 | 379 | 0.7916 | 14987285 | 242 | 0 | 490.22 |
| HEMANGIOL | 5 | 997 | 0.5015 | 14986667 | 240 | 0 | 313.16 |
| HEXYON | 17 | 3945 | 0.4309 | 14983719 | 228 | 0 | 283.2 |
| PREVENAR | 36 | 12728 | 0.2828 | 14974936 | 209 | 0 | 202.66 |
| SYNAGIS | 39 | 16551 | 0.2356 | 14971113 | 206 | 0 | 171.25 |
| INFANRIX | 37 | 21210 | 0.1744 | 14966454 | 208 | 0 | 125.52 |
| ROTATEQ | 19 | 19616 | 0.0969 | 14968048 | 226 | 0 | 64.15 |
| VIRAMUNE | 3 | 3210 | 0.0935 | 14984454 | 242 | 0 | 57.87 |
| SUBOXONE | 7 | 22773 | 0.0307 | 14964891 | 238 | 0 | 19.33 |
| ROTARIX | 8 | 29323 | 0.0273 | 14958341 | 237 | 0 | 17.22 |
| BEXSERO | 10 | 52110 | 0.0192 | 14935554 | 235 | 0 | 12.2 |
| SYNFLORIX | 3 | 15438 | 0.0194 | 14972226 | 242 | 0 | 12.02 |
| KEPPRA | 3 | 38907 | 0.0077 | 14948757 | 242 | 0 | 4.76 |

What is noticeable is the predominance of vaccines in this list (shaded in yellow)

BEYFORTUS : *“Beyfortus is an injection given directly to your baby that provides fast-acting antibody protection (immunization) against serious RSV lung infection” (RSV is respiratory syncytial virus)”*

\$519.75 per dose (highest price for the highest SIDS rate !)

ROTASHIELD : The first rotavirus vaccine, introduced 1998. Caused intussusception and was withdrawn.

HEXYON : *“A hexavalent vaccine, or 6-in-1 vaccine, is a combination vaccine with six individual vaccines conjugated into one, intended to protect people from multiple diseases.”*

“Hexyon (DTaP-IPV-HB-Hib) is indicated for primary and booster vaccination of infants and toddlers from 6 weeks of age against diphtheria, tetanus, pertussis, hepatitis B, poliomyelitis and invasive diseases caused by Haemophilus influenzae type b (Hib)”

PREVENAR : *“Prevenar 13 is used to protect children aged between six weeks and 17 years against invasive disease, pneumonia (infection of the lungs) and acute otitis media (infection of the middle ear) caused by S. pneumoniae.”*

- SYNAGIS : *“SYNAGIS [si-nah-jis] is an FDA-approved prescription injection of antibodies that is given monthly **to help protect high-risk infants from severe RSV disease throughout the RSV season.** Each dose provides protection for about 28–30 days.”*
- INFANRIX : *“**Infanrix hexa** is indicated for primary and booster vaccination of infants from the age of 6 weeks and toddlers against diphtheria, tetanus, pertussis, hepatitis B, poliomyelitis and disease caused by Haemophilus influenzae type b. The use of Infanrix hexa should be in accordance with official recommendations.”*
- ROTATEQ : *“RotaTeq is an oral vaccine used **to help prevent rotavirus infection in children.**”*
- ROTARIX : *“ROTARIX is **a vaccine indicated for the prevention of rotavirus gastroenteritis caused by G1 and non-G1 types (G3, G4, and G9) when administered as a 2-dose series.** ROTARIX is approved for use in infants 6 weeks and up to 24 weeks of age.”*
- BEXSERO : *“Bexsero is a vaccine that protects against invasive meningococcal disease caused by the bacterium Neisseria meningitidis group B”*
- SYNFLORIX : *“**SYNFLORIX™ is a 10-valent pneumococcal conjugate vaccine (PCV-10)** (containing serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F) which has had Marketing Authorisation (MA) in Europe since 2009.”*

EU Safety Signal Tool

You can access this tool here –

<https://knollfrank.github.io/HowBadIsMyBatch/SymptomsCausedByDrugs/?symptom=sudden+infant+death+syndrome>

This database looks at 1504 different drugs, including 100 vaccines. The number shown is the lower confidence limit of the PRR.

Just enter “sudden infant death syndrome” as the symptom.

Select Symptom:

Show entries Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|----------|--|
| HEXYON | 173.18 |
| PREVENAR | 142.35 |
| SYNAGIS | 121.65 |
| INFANRIX | 88.5 |
| ROTATEQ | 40.18 |
| BEXSERO | 6.48 |

Showing 1 to 6 of 6 entries Previous 1 Next

All of the drugs that generate a safety signal for SIDS are vaccines.

INFANTILE APNOEA

Apnoea : *“Apnea is a temporary cessation of breathing. It can refer to a sleep disorder called sleep apnea, or to a condition that affects infants and premature babies”*

| DRUG | Apnoea | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|---------------|--------|------------|--------|-----------|-----------|---------|--------|
| HEMANGIOL | 9 | 997 | 0.9027 | 14986667 | 196 | 0 | 690.23 |
| INFANRIX | 38 | 21210 | 0.1792 | 14966454 | 167 | 0 | 160.56 |
| PREVENAR | 19 | 12728 | 0.1493 | 14974936 | 186 | 0 | 120.18 |
| PROPRANOLOL | 4 | 3922 | 0.102 | 14983742 | 201 | 0 | 76.03 |
| VIRAMUNE | 3 | 3210 | 0.0935 | 14984454 | 202 | 0 | 69.33 |
| HEXYON | 3 | 3945 | 0.076 | 14983719 | 202 | 0 | 56.41 |
| SYNAGIS | 7 | 16551 | 0.0423 | 14971113 | 198 | 0 | 31.98 |
| MIDAZOLAM | 3 | 8126 | 0.0369 | 14979538 | 202 | 0 | 27.38 |
| LEVETIRACETAM | 7 | 24458 | 0.0286 | 14963206 | 198 | 0 | 21.63 |
| ROTATEQ | 5 | 19616 | 0.0255 | 14968048 | 200 | 0 | 19.08 |
| FENTANYL | 7 | 31702 | 0.0221 | 14955962 | 198 | 0 | 16.68 |
| BEXSERO | 11 | 52110 | 0.0211 | 14935554 | 194 | 0 | 16.25 |
| KEPPRA | 6 | 38907 | 0.0154 | 14948757 | 199 | 0 | 11.58 |
| ABILIFY | 3 | 24065 | 0.0125 | 14963599 | 202 | 0 | 9.23 |
| ZYPREXA | 7 | 69954 | 0.01 | 14917710 | 198 | 0 | 7.54 |
| HUMALOG | 4 | 73075 | 0.0055 | 14914589 | 201 | 0 | 4.06 |
| CYMBALTA | 5 | 115064 | 0.0043 | 14872600 | 200 | 0 | 3.23 |

Many of the drugs associated with apnoea are vaccines (shaded in yellow). There are 100 different vaccines in the EU data, but only these 6 vaccines have an association with apnoea. These 6 vaccines appeared before when we looked at SIDS.

Select Symptom:

Show entries Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|----------|--|
| INFANRIX | 112.92 |
| PREVENAR | 74.98 |
| BEXSERO | 8.85 |

Showing 1 to 3 of 3 entries Previous 1 Next

APNOEIC ATTACK

Once again, the criteria applied for a safety signal were symptom ≥ 3 , PRR > 2

| DRUG | apnoeic attack | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|---------------|----------------|------------|--------|-----------|-----------|---------|--------|
| INFANRIX | 33 | 21210 | 0.1556 | 14966454 | 124 | 0 | 187.79 |
| SYNAGIS | 15 | 16551 | 0.0906 | 14971113 | 142 | 0 | 95.55 |
| MIDAZOLAM | 5 | 8126 | 0.0615 | 14979538 | 152 | 0 | 60.64 |
| SYNFLORIX | 5 | 15438 | 0.0324 | 14972226 | 152 | 0 | 31.9 |
| ROTATEQ | 6 | 19616 | 0.0306 | 14968048 | 151 | 0 | 30.32 |
| PREVENAR | 3 | 12728 | 0.0236 | 14974936 | 154 | 0 | 22.92 |
| ROTARIX | 6 | 29323 | 0.0205 | 14958341 | 151 | 0 | 20.27 |
| FENTANYL | 6 | 31702 | 0.0189 | 14955962 | 151 | 0 | 18.75 |
| LEVETIRACETAM | 3 | 24458 | 0.0123 | 14963206 | 154 | 0 | 11.92 |
| BEXSERO | 5 | 52110 | 0.0096 | 14935554 | 152 | 0 | 9.43 |

Select Symptom:

Show entries

Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|----------|--|
| INFANRIX | 127.95 |
| SYNAGIS | 56.13 |

Showing 1 to 2 of 2 entries

Previous 1 Next

APNOEA

This is a general classification, rather than specifically infantile. It shows that dysregulation of the breathing reflex following administration of these drugs occurs regardless of age. You can see that some vaccines shaded in yellow exceed fentanyl in this effect.

| DRUG | apnoea | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|-----------------|--------|------------|--------|-----------|-----------|---------|--------|
| PROCOMVAX | 9 | 157 | 5.7325 | 14987507 | 2163 | 0.01 | 397.21 |
| GENERAL | 3 | 176 | 1.7045 | 14987488 | 2169 | 0.01 | 117.78 |
| NALOXONE | 17 | 1139 | 1.4925 | 14986525 | 2155 | 0.01 | 103.8 |
| SYNAGIS | 197 | 16551 | 1.1903 | 14971113 | 1975 | 0.01 | 90.23 |
| HEXACIMA | 24 | 1976 | 1.2146 | 14985688 | 2148 | 0.01 | 84.74 |
| BEYFORTUS | 5 | 523 | 0.956 | 14987141 | 2167 | 0.01 | 66.12 |
| TRITANRIX | 3 | 326 | 0.9202 | 14987338 | 2169 | 0.01 | 63.59 |
| INFANRIX | 177 | 21210 | 0.8345 | 14966454 | 1995 | 0.01 | 62.6 |
| COLISTIMETHATE | 3 | 427 | 0.7026 | 14987237 | 2169 | 0.01 | 48.55 |
| PREVENAR | 86 | 12728 | 0.6757 | 14974936 | 2086 | 0.01 | 48.51 |
| PECFENT | 3 | 445 | 0.6742 | 14987219 | 2169 | 0.01 | 46.58 |
| SYNFLORIX | 87 | 15438 | 0.5635 | 14972226 | 2085 | 0.01 | 40.47 |
| MIDAZOLAM | 44 | 8126 | 0.5415 | 14979538 | 2128 | 0.01 | 38.12 |
| DEXMEDETOMIDINE | 5 | 1079 | 0.4634 | 14986585 | 2167 | 0.01 | 32.05 |
| SOMATROPIN | 17 | 3852 | 0.4413 | 14983812 | 2155 | 0.01 | 30.69 |
| ROTARIX | 117 | 29323 | 0.399 | 14958341 | 2055 | 0.01 | 29.04 |
| ROTATEQ | 68 | 19616 | 0.3467 | 14968048 | 2104 | 0.01 | 24.66 |
| FENTANYL | 103 | 31702 | 0.3249 | 14955962 | 2069 | 0.01 | 23.49 |
| BUPIVACAINE | 19 | 5783 | 0.3285 | 14981881 | 2153 | 0.01 | 22.86 |
| BRIDION | 13 | 4873 | 0.2668 | 14982791 | 2159 | 0.01 | 18.51 |
| NAGLAZYME | 12 | 4943 | 0.2428 | 14982721 | 2160 | 0.01 | 16.84 |
| HEXYON | 9 | 3945 | 0.2281 | 14983719 | 2163 | 0.01 | 15.8 |
| INVIRASE | 3 | 1317 | 0.2278 | 14986347 | 2169 | 0.01 | 15.74 |
| SPINRAZA | 15 | 6624 | 0.2264 | 14981040 | 2157 | 0.01 | 15.73 |
| LIRAGLUTIDE | 3 | 1519 | 0.1975 | 14986145 | 2169 | 0.01 | 13.65 |
| MYOZYME | 14 | 7717 | 0.1814 | 14979947 | 2158 | 0.01 | 12.59 |
| ALDURAZYME | 6 | 3342 | 0.1795 | 14984322 | 2166 | 0.01 | 12.42 |
| ELAPRASE | 4 | 2270 | 0.1762 | 14985394 | 2168 | 0.01 | 12.18 |
| BENZODIAZEPINE | 4 | 2299 | 0.174 | 14985365 | 2168 | 0.01 | 12.03 |
| SUFENTANIL | 4 | 2572 | 0.1555 | 14985092 | 2168 | 0.01 | 10.75 |
| ZOLGENSMA | 4 | 2796 | 0.1431 | 14984868 | 2168 | 0.01 | 9.89 |
| PROQUAD | 8 | 5666 | 0.1412 | 14981998 | 2164 | 0.01 | 9.78 |
| BEXSERO | 71 | 52110 | 0.1363 | 14935554 | 2101 | 0.01 | 9.69 |
| XYREM | 10 | 8621 | 0.116 | 14979043 | 2162 | 0.01 | 8.04 |
| CEREZYME | 7 | 6185 | 0.1132 | 14981479 | 2165 | 0.01 | 7.83 |
| GLIBENCLAMIDE | 3 | 2774 | 0.1081 | 14984890 | 2169 | 0.01 | 7.47 |
| AREPANRIX | 4 | 3853 | 0.1038 | 14983811 | 2168 | 0.01 | 7.18 |
| PROPRANOLOL | 4 | 3922 | 0.102 | 14983742 | 2168 | 0.01 | 7.05 |
| HBVAXPRO | 3 | 3299 | 0.0909 | 14984365 | 2169 | 0.01 | 6.28 |

| | | | | | | | |
|----------------|----|-------|--------|----------|------|------|------|
| M-M-RVAXPRO | 38 | 42976 | 0.0884 | 14944688 | 2134 | 0.01 | 6.19 |
| TORISEL | 3 | 3449 | 0.087 | 14984215 | 2169 | 0.01 | 6.01 |
| MENVEO | 6 | 7210 | 0.0832 | 14980454 | 2166 | 0.01 | 5.76 |
| BRILIQUE | 9 | 10983 | 0.0819 | 14976681 | 2163 | 0.01 | 5.67 |
| MIRCERA | 6 | 7511 | 0.0799 | 14980153 | 2166 | 0.01 | 5.52 |
| LEVETIRACETAM | 19 | 24458 | 0.0777 | 14963206 | 2153 | 0.01 | 5.4 |
| SONOVUE | 3 | 3950 | 0.0759 | 14983714 | 2169 | 0.01 | 5.25 |
| HYDROCORTISONE | 5 | 7078 | 0.0706 | 14980586 | 2167 | 0.01 | 4.88 |
| DULOXETINE | 7 | 10067 | 0.0695 | 14977597 | 2165 | 0.01 | 4.81 |
| BLINCYTO | 6 | 9491 | 0.0632 | 14978173 | 2166 | 0.01 | 4.37 |
| FYCOMPA | 3 | 4829 | 0.0621 | 14982835 | 2169 | 0.01 | 4.29 |
| IBUPROFEN | 15 | 24346 | 0.0616 | 14963318 | 2157 | 0.01 | 4.27 |
| SIFROL | 3 | 5376 | 0.0558 | 14982288 | 2169 | 0.01 | 3.85 |
| ULTIBRO | 3 | 5417 | 0.0554 | 14982247 | 2169 | 0.01 | 3.83 |
| BUPRENORPHINE | 10 | 19044 | 0.0525 | 14968620 | 2162 | 0.01 | 3.64 |
| LATUDA | 9 | 18492 | 0.0487 | 14969172 | 2163 | 0.01 | 3.37 |
| ASPARAGINASE | 4 | 9516 | 0.042 | 14978148 | 2168 | 0.01 | 2.9 |
| KETEK | 3 | 7257 | 0.0413 | 14980407 | 2169 | 0.01 | 2.86 |
| KEPPRA | 16 | 38907 | 0.0411 | 14948757 | 2156 | 0.01 | 2.85 |
| RELVAR | 3 | 8582 | 0.035 | 14979082 | 2169 | 0.01 | 2.41 |
| OLANZAPINE | 15 | 44801 | 0.0335 | 14942863 | 2157 | 0.01 | 2.32 |
| ZYPREXA | 23 | 69954 | 0.0329 | 14917710 | 2149 | 0.01 | 2.28 |
| ZYPADHERA | 4 | 12269 | 0.0326 | 14975395 | 2168 | 0.01 | 2.25 |
| TRACLEER | 11 | 34525 | 0.0319 | 14953139 | 2161 | 0.01 | 2.2 |
| CERVARIX | 13 | 41519 | 0.0313 | 14946145 | 2159 | 0.01 | 2.17 |
| TWINRIX | 4 | 13417 | 0.0298 | 14974247 | 2168 | 0.01 | 2.06 |
| VIMPAT | 4 | 13790 | 0.029 | 14973874 | 2168 | 0.01 | 2 |

Select Symptom:

Show entries

Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|-----------|--|
| SYNAGIS | <u>78</u> |
| NALOXONE | <u>64.63</u> |
| HEXACIMA | <u>56.81</u> |
| INFANRIX | <u>53.72</u> |
| PREVENAR | <u>39.12</u> |
| SYNFLORIX | <u>32.68</u> |
| MIDAZOLAM | <u>28.3</u> |
| ROTARIX | <u>24.12</u> |
| ROTATEQ | <u>19.38</u> |
| FENTANYL | <u>19.28</u> |

Showing 1 to 10 of 29 entries

BREATH HOLDING

Even when a vaccine does not result in SIDS, you can see that it is associated with an excess incidence of episodes of “breath holding” aka apnoea.

| DRUG | breath holding | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|-----------------|----------------|------------|--------|-----------|-----------|---------|--------|
| SYNFLORIX | 22 | 15438 | 0.1425 | 14972226 | 147 | 0 | 145.14 |
| INFANRIX | 28 | 21210 | 0.132 | 14966454 | 141 | 0 | 140.13 |
| TICAGRELOR | 8 | 9470 | 0.0845 | 14978194 | 161 | 0 | 78.59 |
| PREVENAR | 10 | 12728 | 0.0786 | 14974936 | 159 | 0 | 74 |
| ROTATEQ | 9 | 19616 | 0.0459 | 14968048 | 160 | 0 | 42.92 |
| SYNAGIS | 7 | 16551 | 0.0423 | 14971113 | 162 | 0 | 39.09 |
| M-M- RVAXPRO | 14 | 42976 | 0.0326 | 14944688 | 155 | 0 | 31.41 |
| ROTARIX | 9 | 29323 | 0.0307 | 14958341 | 160 | 0 | 28.69 |
| NIMENRIX | 3 | 10985 | 0.0273 | 14976679 | 166 | 0 | 24.64 |
| BEXSERO | 12 | 52110 | 0.023 | 14935554 | 157 | 0 | 21.91 |
| GARDASIL | 4 | 95990 | 0.0042 | 14891674 | 165 | 0 | 3.76 |

Select Symptom:

Show entries

Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|-----------------|--|
| INFANRIX | 93.43 |
| SYNFLORIX | 92.75 |
| PREVENAR | 39.07 |
| M-M- RVAXPRO | 18.18 |
| BEXSERO | 12.18 |

Showing 1 to 5 of 5 entries

Previous 1 Next

POOR FEEDING INFANT

I am not sure why appetite is suppressed. It could be either due to discomfort caused by headaches or nausea, or to dysregulation of the hunger reflex. Later I will show that the thirst reflex is also affected.

| DRUG | poor feeding infant | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|---------------|---------------------|------------|--------|-----------|-----------|---------|--------|
| ROTARIX | 93 | 29323 | 0.3172 | 14958341 | 463 | 0 | 102.47 |
| PREVENAR | 39 | 12728 | 0.3064 | 14974936 | 517 | 0 | 88.75 |
| SYNAGIS | 33 | 16551 | 0.1994 | 14971113 | 523 | 0 | 57.07 |
| BUVIDAL | 3 | 1699 | 0.1766 | 14985965 | 553 | 0 | 47.85 |
| BEXSERO | 66 | 52110 | 0.1267 | 14935554 | 490 | 0 | 38.61 |
| INFANRIX | 26 | 21210 | 0.1226 | 14966454 | 530 | 0 | 34.62 |
| ROTATEQ | 23 | 19616 | 0.1173 | 14968048 | 533 | 0 | 32.93 |
| SUBOXONE | 19 | 22773 | 0.0834 | 14964891 | 537 | 0 | 23.25 |
| BUPRENORPHINE | 15 | 19044 | 0.0788 | 14968620 | 541 | 0 | 21.79 |
| SYNFLORIX | 9 | 15438 | 0.0583 | 14972226 | 547 | 0 | 15.96 |
| MIDAZOLAM | 3 | 8126 | 0.0369 | 14979538 | 553 | 0 | 10 |
| ABILIFY | 5 | 24065 | 0.0208 | 14963599 | 551 | 0 | 5.64 |
| LEVETIRACETAM | 5 | 24458 | 0.0204 | 14963206 | 551 | 0 | 5.55 |
| ARIPIPRAZOLE | 3 | 16231 | 0.0185 | 14971433 | 553 | 0 | 5 |
| KEPPRA | 7 | 38907 | 0.018 | 14948757 | 549 | 0 | 4.9 |
| PANDEMRIX | 8 | 50551 | 0.0158 | 14937113 | 548 | 0 | 4.31 |
| ZYPREXA | 11 | 69954 | 0.0157 | 14917710 | 545 | 0 | 4.3 |
| OLANZAPINE | 6 | 44801 | 0.0134 | 14942863 | 550 | 0 | 3.64 |
| FENTANYL | 3 | 31702 | 0.0095 | 14955962 | 553 | 0 | 2.56 |
| HUMALOG | 6 | 73075 | 0.0082 | 14914589 | 550 | 0 | 2.23 |

FLOPPY INFANT

“Infants with hypotonia may have poor head control, seem floppy, and rest with their elbows and knees loosely extended. They may also have delayed motor skills and poor control of movement.”

| DRUG | floppy infant | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|----------|---------------|------------|--------|-----------|-----------|---------|--------|
| BEXSERO | 38 | 52110 | 0.0729 | 14935554 | 59 | 0 | 184.6 |
| ROTARIX | 19 | 29323 | 0.0648 | 14958341 | 78 | 0 | 124.26 |
| INFANRIX | 13 | 21210 | 0.0613 | 14966454 | 84 | 0 | 109.2 |
| PREVENAR | 5 | 12728 | 0.0393 | 14974936 | 92 | 0 | 63.94 |
| ROTATEQ | 3 | 19616 | 0.0153 | 14968048 | 94 | 0 | 24.35 |

Considering that this database looks at 1504 different drugs, it is extraordinary that the list of vaccines appearing here is the same as appeared for SIDS and for APNOEA.

Select Symptom: floppy infant

Show 10 entries Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|----------|--|
| BEXSERO | 122.81 |
| ROTARIX | 75.27 |
| INFANRIX | 60.9 |

Showing 1 to 3 of 3 entries

Previous 1 Next

INFANTILE SPASMS

| DRUG | infantile spasms | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|---------------|------------------|------------|--------|-----------|-----------|---------|--------|
| VIGABATRIN | 10 | 1302 | 0.768 | 14986362 | 127 | 0 | 906.32 |
| HEXYON | 4 | 3945 | 0.1014 | 14983719 | 133 | 0 | 114.23 |
| INFANRIX | 19 | 21210 | 0.0896 | 14966454 | 118 | 0 | 113.62 |
| EVEROLIMUS | 4 | 4206 | 0.0951 | 14983458 | 133 | 0 | 107.14 |
| PREVENAR | 10 | 12728 | 0.0786 | 14974936 | 127 | 0 | 92.64 |
| SYNAGIS | 12 | 16551 | 0.0725 | 14971113 | 125 | 0 | 86.84 |
| LEVETIRACETAM | 17 | 24458 | 0.0695 | 14963206 | 120 | 0 | 86.67 |
| ROTATEQ | 10 | 19616 | 0.051 | 14968048 | 127 | 0 | 60.08 |
| BEXSERO | 13 | 52110 | 0.0249 | 14935554 | 124 | 0 | 30.05 |
| ROTARIX | 7 | 29323 | 0.0239 | 14958341 | 130 | 0 | 27.47 |
| KEPPRA | 6 | 38907 | 0.0154 | 14948757 | 131 | 0 | 17.6 |
| M-M-RVAXPRO | 3 | 42976 | 0.007 | 14944688 | 134 | 0 | 7.79 |
| PANDEMRIX | 3 | 50551 | 0.0059 | 14937113 | 134 | 0 | 6.62 |

Select Symptom: infantile spasms

Show 10 entries infantile spasms Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|---------------|--|
| VIGABATRIN | 477.2 |
| INFANRIX | 70 |
| LEVETIRACETAM | 52.16 |
| PREVENAR | 48.68 |
| SYNAGIS | 48.03 |
| ROTATEQ | 31.57 |
| BEXSERO | 16.97 |

Showing 1 to 7 of 7 entries Previous 1 Next

Notice the same list – Infanrix, Prevenar, Synagis, Rotateq, Bexsero.

OPISTHOTONUS

Opisthotonus is a condition that causes a person to arch their back and neck into a rigid, abnormal position.

A person with opisthotonus will have a severe spasm of the muscles in their neck, back, and trunk, causing them to arch backward. If they are on their back, only their heels and the back of their head will touch the surface. If they are on their side, their body may resemble a crescent moon.

| DRUG | opisthotonus | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|-----------------|--------------|------------|--------|-----------|-----------|---------|--------|
| HEXYON | 14 | 3945 | 0.3549 | 14983719 | 206 | 0 | 258.13 |
| HEXACIMA | 5 | 1976 | 0.253 | 14985688 | 215 | 0 | 176.37 |
| INFANRIX | 28 | 21210 | 0.132 | 14966454 | 192 | 0 | 102.9 |
| SYNFLORIX | 14 | 15438 | 0.0907 | 14972226 | 206 | 0 | 65.91 |
| ROTARIX | 14 | 29323 | 0.0477 | 14958341 | 206 | 0 | 34.67 |
| MIDAZOLAM | 4 | 8126 | 0.0492 | 14979538 | 216 | 0 | 34.14 |
| PREVENAR | 5 | 12728 | 0.0393 | 14974936 | 215 | 0 | 27.36 |
| NIMENRIX | 4 | 10985 | 0.0364 | 14976679 | 216 | 0 | 25.25 |
| ROTATEQ | 7 | 19616 | 0.0357 | 14968048 | 213 | 0 | 25.08 |
| BEXSERO | 17 | 52110 | 0.0326 | 14935554 | 203 | 0 | 24 |
| M-M- RVAXPRO | 10 | 42976 | 0.0233 | 14944688 | 210 | 0 | 16.56 |
| GARDASIL9 | 8 | 41282 | 0.0194 | 14946382 | 212 | 0 | 13.66 |
| ZYPREXA | 13 | 69954 | 0.0186 | 14917710 | 207 | 0 | 13.39 |
| FENTANYL | 4 | 31702 | 0.0126 | 14955962 | 216 | 0 | 8.74 |
| OLANZAPINE | 5 | 44801 | 0.0112 | 14942863 | 215 | 0 | 7.76 |
| GARDASIL | 5 | 95990 | 0.0052 | 14891674 | 215 | 0 | 3.61 |

HYPOTONIC-HYPORESPONSIVE EPISODE

| DRUG | hypotonic-hyporesponsive episode | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|--------------|----------------------------------|------------|--------|-----------|-----------|---------|--------|
| SYNFLORIX | 507 | 15438 | 3.2841 | 14972226 | 1890 | 0.01 | 260.16 |
| TRIACELLUVAX | 4 | 101 | 3.9604 | 14987563 | 2393 | 0.02 | 248.04 |
| HEXACIMA | 65 | 1976 | 3.2895 | 14985688 | 2332 | 0.02 | 211.39 |
| INFANRIX | 506 | 21210 | 2.3857 | 14966454 | 1891 | 0.01 | 188.82 |
| HEXYON | 45 | 3945 | 1.1407 | 14983719 | 2352 | 0.02 | 72.67 |
| AREPANRIX | 35 | 3853 | 0.9084 | 14983811 | 2362 | 0.02 | 57.63 |
| ROTARIX | 225 | 29323 | 0.7673 | 14958341 | 2172 | 0.01 | 52.84 |
| PREVENAR | 91 | 12728 | 0.715 | 14974936 | 2306 | 0.02 | 46.43 |
| BEXSERO | 193 | 52110 | 0.3704 | 14935554 | 2204 | 0.01 | 25.1 |
| SILGARD | 4 | 1072 | 0.3731 | 14986592 | 2393 | 0.02 | 23.37 |
| M-M-RVAXPRO | 115 | 42976 | 0.2676 | 14944688 | 2282 | 0.02 | 17.52 |
| ROTATEQ | 50 | 19616 | 0.2549 | 14968048 | 2347 | 0.02 | 16.26 |
| NOT | 23 | 9102 | 0.2527 | 14978562 | 2374 | 0.02 | 15.94 |
| HBVAXPRO | 6 | 3299 | 0.1819 | 14984365 | 2391 | 0.02 | 11.4 |
| NIMENRIX | 10 | 10985 | 0.091 | 14976679 | 2387 | 0.02 | 5.71 |
| PROQUAD | 5 | 5666 | 0.0882 | 14981998 | 2392 | 0.02 | 5.53 |
| TWINRIX | 8 | 13417 | 0.0596 | 14974247 | 2389 | 0.02 | 3.74 |

Select Symptom: hypotonic-hyporesponsive episode

Show 10 entries

Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|-------------|--|
| SYNFLORIX | 236.17 |
| INFANRIX | 171.33 |
| HEXACIMA | 165.87 |
| HEXYON | 54.2 |
| ROTARIX | 46.09 |
| AREPANRIX | 41.33 |
| PREVENAR | 37.68 |
| BEXSERO | 21.67 |
| M-M-RVAXPRO | 14.53 |
| ROTATEQ | 12.29 |

Showing 1 to 10 of 12 entries

Previous 1 2 Next

HYPOTONIA

Notice that the incidence of hypotonia with the shaded vaccines far exceeds that of cannabidiol.

| DRUG | hypotonia | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|-----------------|-----------|------------|--------|-----------|-----------|---------|--------|
| TRITANRIX | 15 | 326 | 4.6012 | 14987338 | 6162 | 0.04 | 111.91 |
| POLIOMYELITIS | 5 | 141 | 3.5461 | 14987523 | 6172 | 0.04 | 86.11 |
| DIPHTHERIA | 4 | 116 | 3.4483 | 14987548 | 6173 | 0.04 | 83.72 |
| ROTASHIELD | 6 | 379 | 1.5831 | 14987285 | 6171 | 0.04 | 38.45 |
| HEXAVAC | 3 | 194 | 1.5464 | 14987470 | 6174 | 0.04 | 37.54 |
| HEXACIMA | 29 | 1976 | 1.4676 | 14985688 | 6148 | 0.04 | 35.77 |
| HEXYON | 57 | 3945 | 1.4449 | 14983719 | 6120 | 0.04 | 35.37 |
| INFANRIX | 250 | 21210 | 1.1787 | 14966454 | 5927 | 0.04 | 29.76 |
| NON | 3 | 257 | 1.1673 | 14987407 | 6174 | 0.04 | 28.34 |
| SYNFLORIX | 121 | 15438 | 0.7838 | 14972226 | 6056 | 0.04 | 19.38 |
| PREVENAR | 96 | 12728 | 0.7542 | 14974936 | 6081 | 0.04 | 18.57 |
| BEXSERO | 371 | 52110 | 0.712 | 14935554 | 5806 | 0.04 | 18.31 |
| ODOMZO | 6 | 916 | 0.655 | 14986748 | 6171 | 0.04 | 15.91 |
| ROTARIX | 163 | 29323 | 0.5559 | 14958341 | 6014 | 0.04 | 13.83 |
| DIACOMIT | 13 | 2321 | 0.5601 | 14985343 | 6164 | 0.04 | 13.62 |
| DEXMEDETOMIDINE | 6 | 1079 | 0.5561 | 14986585 | 6171 | 0.04 | 13.5 |
| MYOZYME | 41 | 7717 | 0.5313 | 14979947 | 6136 | 0.04 | 12.97 |
| BRINEURA | 4 | 778 | 0.5141 | 14986886 | 6173 | 0.04 | 12.48 |
| DEXDOR | 3 | 629 | 0.4769 | 14987035 | 6174 | 0.04 | 11.58 |
| ROTATEQ | 76 | 19616 | 0.3874 | 14968048 | 6101 | 0.04 | 9.51 |
| PROQUAD | 22 | 5666 | 0.3883 | 14981998 | 6155 | 0.04 | 9.45 |
| CELVAPAN | 6 | 1897 | 0.3163 | 14985767 | 6171 | 0.04 | 7.68 |
| INVIRASE | 4 | 1317 | 0.3037 | 14986347 | 6173 | 0.04 | 7.37 |
| NOT | 27 | 9102 | 0.2966 | 14978562 | 6150 | 0.04 | 7.22 |
| ZOLGENSMA | 7 | 2796 | 0.2504 | 14984868 | 6170 | 0.04 | 6.08 |
| HBVAXPRO | 8 | 3299 | 0.2425 | 14984365 | 6169 | 0.04 | 5.89 |
| IXIARO | 6 | 2603 | 0.2305 | 14985061 | 6171 | 0.04 | 5.6 |
| VIGABATRIN | 3 | 1302 | 0.2304 | 14986362 | 6174 | 0.04 | 5.59 |
| JCOVDEN | 4 | 1763 | 0.2269 | 14985901 | 6173 | 0.04 | 5.51 |
| SYNAGIS | 37 | 16551 | 0.2236 | 14971113 | 6140 | 0.04 | 5.45 |
| MENVEO | 15 | 7210 | 0.208 | 14980454 | 6162 | 0.04 | 5.06 |
| FYCOMPA | 10 | 4829 | 0.2071 | 14982835 | 6167 | 0.04 | 5.03 |
| PANDEMRIX | 103 | 50551 | 0.2038 | 14937113 | 6074 | 0.04 | 5.01 |
| M-M-RVAXPRO | 86 | 42976 | 0.2001 | 14944688 | 6091 | 0.04 | 4.91 |
| SUFENTANIL | 5 | 2572 | 0.1944 | 14985092 | 6172 | 0.04 | 4.72 |
| CANNABIDIOL | 4 | 2070 | 0.1932 | 14985594 | 6173 | 0.04 | 4.69 |
| STRENSIQ | 4 | 2206 | 0.1813 | 14985458 | 6173 | 0.04 | 4.4 |
| UROREC | 3 | 1671 | 0.1795 | 14985993 | 6174 | 0.04 | 4.36 |
| PROPRANOLOL | 7 | 3922 | 0.1785 | 14983742 | 6170 | 0.04 | 4.33 |
| EBIXA | 5 | 2845 | 0.1757 | 14984819 | 6172 | 0.04 | 4.27 |

| | | | | | | | |
|----------------|-----|--------|--------|----------|------|------|------|
| ZYPADHERA | 21 | 12269 | 0.1712 | 14975395 | 6156 | 0.04 | 4.16 |
| SEBIVO | 5 | 2938 | 0.1702 | 14984726 | 6172 | 0.04 | 4.13 |
| MEMANTINE | 4 | 2415 | 0.1656 | 14985249 | 6173 | 0.04 | 4.02 |
| BRIDION | 8 | 4873 | 0.1642 | 14982791 | 6169 | 0.04 | 3.99 |
| VALSARTAN | 9 | 5641 | 0.1595 | 14982023 | 6168 | 0.04 | 3.88 |
| NILEMDO | 5 | 3302 | 0.1514 | 14984362 | 6172 | 0.04 | 3.68 |
| NAGLAZYME | 7 | 4943 | 0.1416 | 14982721 | 6170 | 0.04 | 3.44 |
| SPINRAZA | 9 | 6624 | 0.1359 | 14981040 | 6168 | 0.04 | 3.3 |
| LEVETIRACETAM | 33 | 24458 | 0.1349 | 14963206 | 6144 | 0.04 | 3.29 |
| COMBIVIR | 7 | 5212 | 0.1343 | 14982452 | 6170 | 0.04 | 3.26 |
| BENZODIAZEPINE | 3 | 2299 | 0.1305 | 14985365 | 6174 | 0.04 | 3.17 |
| AREPANRIX | 5 | 3853 | 0.1298 | 14983811 | 6172 | 0.04 | 3.15 |
| CERVARIX | 53 | 41519 | 0.1277 | 14946145 | 6124 | 0.04 | 3.12 |
| TOPOTECAN | 4 | 3129 | 0.1278 | 14984535 | 6173 | 0.04 | 3.1 |
| GILENYA | 3 | 2356 | 0.1273 | 14985308 | 6174 | 0.04 | 3.09 |
| MIDAZOLAM | 10 | 8126 | 0.1231 | 14979538 | 6167 | 0.04 | 2.99 |
| FERRIPROX | 3 | 2474 | 0.1213 | 14985190 | 6174 | 0.04 | 2.94 |
| NIMENRIX | 13 | 10985 | 0.1183 | 14976679 | 6164 | 0.04 | 2.88 |
| LEVODOPA | 3 | 2559 | 0.1172 | 14985105 | 6174 | 0.04 | 2.85 |
| FLUENZ | 11 | 9446 | 0.1165 | 14978218 | 6166 | 0.04 | 2.83 |
| SIMBRINZA | 4 | 3444 | 0.1161 | 14984220 | 6173 | 0.04 | 2.82 |
| VAXZEVRIA | 775 | 746718 | 0.1038 | 14240946 | 5402 | 0.04 | 2.74 |
| GARDASIL9 | 44 | 41282 | 0.1066 | 14946382 | 6133 | 0.04 | 2.6 |
| TWINRIX | 14 | 13417 | 0.1043 | 14974247 | 6163 | 0.04 | 2.54 |
| EPIVIR | 4 | 3842 | 0.1041 | 14983822 | 6173 | 0.04 | 2.53 |
| KEPPRA | 38 | 38907 | 0.0977 | 14948757 | 6139 | 0.04 | 2.38 |
| PRIALT | 4 | 4111 | 0.0973 | 14983553 | 6173 | 0.04 | 2.36 |
| FOCETRIA | 7 | 7862 | 0.089 | 14979802 | 6170 | 0.04 | 2.16 |
| GARDASIL | 81 | 95990 | 0.0844 | 14891674 | 6096 | 0.04 | 2.06 |
| SELINCRO | 3 | 3580 | 0.0838 | 14984084 | 6174 | 0.04 | 2.03 |
| DENGVAXIA | 4 | 4844 | 0.0826 | 14982820 | 6173 | 0.04 | 2 |

Select Symptom:

Show entries

Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|-----------|--|
| TRITANRIX | 68.22 |
| HEXYON | 27.3 |
| INFANRIX | 26.24 |
| HEXACIMA | 24.9 |
| BEXSERO | 16.5 |
| SYNFLORIX | 16.2 |
| PREVENAR | 15.19 |
| ROTARIX | 11.84 |
| MYOZYME | 9.55 |
| DIACOMIT | 7.91 |

Showing 1 to 10 of 32 entries

[Previous](#) [1](#) [2](#) [3](#) [4](#) [Next](#)

FONTANELLE BULGING

A bulging fontanelle is a medical emergency that occurs when an infant's soft spot curves outward due to fluid buildup or increased pressure in the brain. It can be a sign of a serious underlying condition, such as meningitis, encephalitis, or hydrocephalus.

| DRUG | fontanelle bulging | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|-----------|--------------------|------------|--------|-----------|-----------|---------|--------|
| PREVENAR | 12 | 12728 | 0.0943 | 14974936 | 84 | 0 | 168.08 |
| SYNAGIS | 11 | 16551 | 0.0665 | 14971113 | 85 | 0 | 117.06 |
| INFANRIX | 12 | 21210 | 0.0566 | 14966454 | 84 | 0 | 100.8 |
| ROTATEQ | 10 | 19616 | 0.051 | 14968048 | 86 | 0 | 88.73 |
| BEXSERO | 18 | 52110 | 0.0345 | 14935554 | 78 | 0 | 66.14 |
| SYNFLORIX | 6 | 15438 | 0.0389 | 14972226 | 90 | 0 | 64.66 |
| ROTARIX | 7 | 29323 | 0.0239 | 14958341 | 89 | 0 | 40.12 |

Select Symptom: fontanelle bulging

Show 10 entries

Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|----------|--|
| PREVENAR | 91.82 |
| SYNAGIS | 62.48 |
| INFANRIX | 55.06 |
| ROTATEQ | 46.1 |
| BEXSERO | 39.62 |

Showing 1 to 5 of 5 entries

Previous 1 Next

OLIGODIPSIA

Oligodipsia is a condition where a person has little or no thirst.

| DRUG | oligodipsia | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|-------------|-------------|------------|--------|-----------|-----------|---------|--------|
| INFANRIX | 31 | 21210 | 0.1462 | 14966454 | 113 | 0 | 193.58 |
| PANDEMRIX | 41 | 50551 | 0.0811 | 14937113 | 103 | 0 | 117.62 |
| AXITINIB | 3 | 4092 | 0.0733 | 14983572 | 141 | 0 | 77.91 |
| M-M-RVAXPRO | 15 | 42976 | 0.0349 | 14944688 | 129 | 0 | 40.44 |
| ROTARIX | 7 | 29323 | 0.0239 | 14958341 | 137 | 0 | 26.06 |

Select Symptom:

Show entries

Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|-------------|--|
| INFANRIX | 130.13 |
| PANDEMRIX | 81.91 |
| M-M-RVAXPRO | 23.69 |

Showing 1 to 3 of 3 entries

Previous 1 Next

Taking a Closer Look at Infanrix

This data is available with the EU Safety Signal :

<https://knollfrank.github.io/HowBadIsMyBatch/SymptomsCausedByDrugs/?symptom=infantile+spasms&vaccine=INFANRIX>

Strongest Symptoms for "INFANRIX"

| Symptom | Lower Confidence Limit of Proportional Reporting Ratio ≥ 2 |
|----------------------------------|---|
| high-pitched crying | 173.85 |
| hypotonic-hyporesponsive episode | 171.33 |
| brief resolved unexplained event | 136.68 |
| oligodipsia | 130.13 |
| apnoeic attack | 127.95 |
| infantile apnoea | 112.92 |
| crying | 106.62 |
| breath holding | 93.43 |
| sudden infant death syndrome | 88.5 |
| infantile spasms | 70 |
| opisthotonus | 69.24 |
| decreased eye contact | 65.3 |
| kawasaki's disease | 62.54 |
| psychomotor skills impaired | 62.17 |
| floppy infant | 60.9 |
| fontanelle bulging | 55.06 |
| screaming | 54.47 |
| apnoea | 53.72 |
| febrile convulsion | 53.34 |
| developmental delay | 51.76 |
| speech disorder developmental | 43.15 |
| autism spectrum disorder | 35.53 |
| staring | 34.11 |
| hyporesponsive to stimuli | 33.85 |
| moaning | 32.72 |
| vaccination site granuloma | 31.79 |
| hypertonia | 28.83 |

There are many symptoms that are related to SIDS –

- Hypotonic-hyporesponsive episode
- Apnoeic attack
- Infantile apnoea
- Breath holding
- Sudden Infant Death Syndrome
- Infantile spasms
- Floppy Infant
- Apnoea
- Hypotonia

Given the association of SIDS with APNOEA , HYPOTONIA, and OLIGODYPSIA it seems that a contributing factor to sudden infant death is the effect of vaccines on the breathing reflex – as seen in apnoea and hyotonia.

This, in turn, may be related to inflammation of the brain – visible as fontanelle bulging – and further indicated by spasms and convulsions, and by various symptoms of autism.

“BRIEF RESOLVED UNEXPLAINED EVENT”

In the symptom list above for INFANRIX, you might have noticed an odd entry – BRIEF RESOLVED UNEXPLAINED EVENT.

We might ask if such events are only associated with certain drugs ?

| DRUG | brief resolved unexplained event | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|-----------|----------------------------------|------------|--------|-----------|-----------|---------|--------|
| INFANRIX | 29 | 21210 | 0.1367 | 14966454 | 99 | 0 | 206.7 |
| PREVENAR | 13 | 12728 | 0.1021 | 14974936 | 115 | 0 | 133 |
| HEXYON | 4 | 3945 | 0.1014 | 14983719 | 124 | 0 | 122.52 |
| ROTARIX | 14 | 29323 | 0.0477 | 14958341 | 114 | 0 | 62.65 |
| SYNFLORIX | 7 | 15438 | 0.0453 | 14972226 | 121 | 0 | 56.11 |
| ROTATEQ | 6 | 19616 | 0.0306 | 14968048 | 122 | 0 | 37.53 |
| SYNAGIS | 5 | 16551 | 0.0302 | 14971113 | 123 | 0 | 36.77 |
| BEXSERO | 13 | 52110 | 0.0249 | 14935554 | 115 | 0 | 32.4 |

Oddly enough, it is the same list of nasties we have come to recognize.

It seems that what is “unexplained” might be explained through its particular association with certain drugs.

Select Symptom:

Show entries Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|----------|--|
| INFANRIX | 136.68 |
| PREVENAR | 74.97 |
| ROTARIX | 35.97 |
| BEXSERO | 18.26 |

Showing 1 to 4 of 4 entries Previous 1 Next

HIGH PITCHED CRYING

Another odd symptom in the list is HIGH PITCHED CRYING – which sounds a bit like screaming. Rather than expressing mild discomfort – it expresses agony.

| DRUG | high-pitched crying | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|-------------|---------------------|------------|--------|-----------|-----------|---------|--------|
| INFANRIX | 47 | 21210 | 0.2216 | 14966454 | 137 | 0 | 242.08 |
| HEXYON | 11 | 3945 | 0.2788 | 14983719 | 173 | 0 | 241.5 |
| HEXACIMA | 3 | 1976 | 0.1518 | 14985688 | 181 | 0 | 125.7 |
| BEXSERO | 44 | 52110 | 0.0844 | 14935554 | 140 | 0 | 90.08 |
| SYNFLORIX | 10 | 15438 | 0.0648 | 14972226 | 174 | 0 | 55.74 |
| PREVENAR | 8 | 12728 | 0.0629 | 14974936 | 176 | 0 | 53.48 |
| PROQUAD | 3 | 5666 | 0.0529 | 14981998 | 181 | 0 | 43.83 |
| ROTARIX | 14 | 29323 | 0.0477 | 14958341 | 170 | 0 | 42.01 |
| ROTATEQ | 9 | 19616 | 0.0459 | 14968048 | 175 | 0 | 39.24 |
| M-M-RVAXPRO | 4 | 42976 | 0.0093 | 14944688 | 180 | 0 | 7.73 |

Select Symptom:

Show entries

Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|-----------|--|
| INFANRIX | 173.85 |
| HEXYON | 131.4 |
| BEXSERO | 64.2 |
| SYNFLORIX | 29.47 |
| ROTARIX | 24.36 |

Showing 1 to 5 of 5 entries

Previous 1 Next

CRYING

Given that crying is an index of discomfort, lets look at all 1504 drugs in the Eudra dataset to see if there are any drugs that stand out for causing crying.

You can see that INFANRIX occasions 112 x the average incidence of crying. Perhaps our infants are trying to tell us something.

The highest incidence of CRYING occurs with the same drugs that exhibit the highest incidence of HIGH PITCHED CRYING. Remember that this is comparing all 1504 drugs, so it is remarkable that these same drugs stand out.

| DRUG | crying | DRUG-TOTAL | % | OTHER-TOT | OTHER-SYM | % OTHER | PRR |
|--------------|--------|------------|--------|-----------|-----------|---------|--------|
| HEXACIMA | 166 | 1976 | 8.4008 | 14985688 | 10438 | 0.07 | 120.61 |
| INFANRIX | 1458 | 21210 | 6.8741 | 14966454 | 9146 | 0.06 | 112.49 |
| SYNFLORIX | 1066 | 15438 | 6.905 | 14972226 | 9538 | 0.06 | 108.39 |
| TRITANRIX | 20 | 326 | 6.135 | 14987338 | 10584 | 0.07 | 86.87 |
| PROCOMVAX | 6 | 157 | 3.8217 | 14987507 | 10598 | 0.07 | 54.05 |
| HEXYON | 119 | 3945 | 3.0165 | 14983719 | 10485 | 0.07 | 43.11 |
| ALLI | 10 | 348 | 2.8736 | 14987316 | 10594 | 0.07 | 40.65 |
| ROTARIX | 755 | 29323 | 2.5748 | 14958341 | 9849 | 0.07 | 39.1 |
| DIPHThERIA | 3 | 116 | 2.5862 | 14987548 | 10601 | 0.07 | 36.56 |
| ROTATEQ | 457 | 19616 | 2.3297 | 14968048 | 10147 | 0.07 | 34.37 |
| PREVENAR | 296 | 12728 | 2.3256 | 14974936 | 10308 | 0.07 | 33.78 |
| M-M-RVAXPRO | 897 | 42976 | 2.0872 | 14944688 | 9707 | 0.06 | 32.13 |
| BEXSERO | 1032 | 52110 | 1.9804 | 14935554 | 9572 | 0.06 | 30.9 |
| NUEDEXTA | 5 | 314 | 1.5924 | 14987350 | 10599 | 0.07 | 22.52 |
| HEXAVAC | 3 | 194 | 1.5464 | 14987470 | 10601 | 0.07 | 21.86 |
| NGENLA | 7 | 603 | 1.1609 | 14987061 | 10597 | 0.07 | 16.42 |
| SLENYTO | 5 | 495 | 1.0101 | 14987169 | 10599 | 0.07 | 14.28 |
| ACOMPLIA | 50 | 5238 | 0.9546 | 14982426 | 10554 | 0.07 | 13.55 |
| HEMANGIOL | 9 | 997 | 0.9027 | 14986667 | 10595 | 0.07 | 12.77 |
| PROQUAD | 48 | 5666 | 0.8472 | 14981998 | 10556 | 0.07 | 12.02 |
| ROTASHIELD | 3 | 379 | 0.7916 | 14987285 | 10601 | 0.07 | 11.19 |
| BREXPIRAZOLE | 3 | 383 | 0.7833 | 14987281 | 10601 | 0.07 | 11.07 |
| SYNAGIS | 126 | 16551 | 0.7613 | 14971113 | 10478 | 0.07 | 10.88 |
| MENQUADFI | 4 | 532 | 0.7519 | 14987132 | 10600 | 0.07 | 10.63 |
| ZANAMIVIR | 4 | 675 | 0.5926 | 14986989 | 10600 | 0.07 | 8.38 |
| HBVAXPRO | 18 | 3299 | 0.5456 | 14984365 | 10586 | 0.07 | 7.72 |
| CHAMPIX | 223 | 41627 | 0.5357 | 14946037 | 10381 | 0.07 | 7.71 |
| RIMONABANT | 5 | 1027 | 0.4869 | 14986637 | 10599 | 0.07 | 6.88 |
| NIMENRIX | 52 | 10985 | 0.4734 | 14976679 | 10552 | 0.07 | 6.72 |
| LATUDA | 73 | 18492 | 0.3948 | 14969172 | 10531 | 0.07 | 5.61 |
| TECFIDERA | 5 | 1404 | 0.3561 | 14986260 | 10599 | 0.07 | 5.04 |
| PANDEMRIX | 175 | 50551 | 0.3462 | 14937113 | 10429 | 0.07 | 4.96 |
| RESOLOR | 5 | 1537 | 0.3253 | 14986127 | 10599 | 0.07 | 4.6 |

| | | | | | | | |
|---------------|-----|--------|--------|----------|-------|------|------|
| BRINTELLIX | 48 | 14994 | 0.3201 | 14972670 | 10556 | 0.07 | 4.54 |
| VIGABATRIN | 4 | 1302 | 0.3072 | 14986362 | 10600 | 0.07 | 4.34 |
| VARENICLINE | 5 | 1671 | 0.2992 | 14985993 | 10599 | 0.07 | 4.23 |
| VORTIOXETINE | 3 | 1072 | 0.2799 | 14986592 | 10601 | 0.07 | 3.96 |
| CYMBALTA | 310 | 115064 | 0.2694 | 14872600 | 10294 | 0.07 | 3.89 |
| REBIF | 4 | 1743 | 0.2295 | 14985921 | 10600 | 0.07 | 3.24 |
| ELAPRASE | 5 | 2270 | 0.2203 | 14985394 | 10599 | 0.07 | 3.11 |
| VALDOXAN | 21 | 9635 | 0.218 | 14978029 | 10583 | 0.07 | 3.08 |
| SPRAVATO | 9 | 4200 | 0.2143 | 14983464 | 10595 | 0.07 | 3.03 |
| AREPANRIX | 8 | 3853 | 0.2076 | 14983811 | 10596 | 0.07 | 2.94 |
| ZOELY | 6 | 2917 | 0.2057 | 14984747 | 10598 | 0.07 | 2.91 |
| MIRAPEXIN | 3 | 1474 | 0.2035 | 14986190 | 10601 | 0.07 | 2.88 |
| TRAVATAN | 4 | 2006 | 0.1994 | 14985658 | 10600 | 0.07 | 2.82 |
| AERIUS | 12 | 6204 | 0.1934 | 14981460 | 10592 | 0.07 | 2.74 |
| UNSPECIFIED | 4 | 2080 | 0.1923 | 14985584 | 10600 | 0.07 | 2.72 |
| DALIRESP | 5 | 2606 | 0.1919 | 14985058 | 10599 | 0.07 | 2.71 |
| TAMIFLU | 28 | 15594 | 0.1796 | 14972070 | 10576 | 0.07 | 2.54 |
| ZOLGENSMA | 5 | 2796 | 0.1788 | 14984868 | 10599 | 0.07 | 2.53 |
| OMNITROPE | 3 | 1688 | 0.1777 | 14985976 | 10601 | 0.07 | 2.51 |
| BRIVIACT | 8 | 4541 | 0.1762 | 14983123 | 10596 | 0.07 | 2.49 |
| SUBOXONE | 40 | 22773 | 0.1756 | 14964891 | 10564 | 0.07 | 2.49 |
| OCRELIZUMAB | 4 | 2283 | 0.1752 | 14985381 | 10600 | 0.07 | 2.48 |
| INTUNIV | 10 | 5823 | 0.1717 | 14981841 | 10594 | 0.07 | 2.43 |
| MENVEO | 12 | 7210 | 0.1664 | 14980454 | 10592 | 0.07 | 2.35 |
| FYCOMPA | 8 | 4829 | 0.1657 | 14982835 | 10596 | 0.07 | 2.34 |
| MEMANTINE | 4 | 2415 | 0.1656 | 14985249 | 10600 | 0.07 | 2.34 |
| NAGLAZYME | 8 | 4943 | 0.1618 | 14982721 | 10596 | 0.07 | 2.29 |
| EXTAVIA | 18 | 11364 | 0.1584 | 14976300 | 10586 | 0.07 | 2.24 |
| DESLORATADINE | 4 | 2569 | 0.1557 | 14985095 | 10600 | 0.07 | 2.2 |
| FLUENZ | 14 | 9446 | 0.1482 | 14978218 | 10590 | 0.07 | 2.1 |
| LYRICA | 42 | 28323 | 0.1483 | 14959341 | 10562 | 0.07 | 2.1 |
| TIVICAY | 5 | 3433 | 0.1456 | 14984231 | 10599 | 0.07 | 2.06 |

Select Symptom:

Show entries

Search:

| Drug | Lower Confidence Limit of Proportional Reporting Ratio |
|-----------------|--|
| INFANRIX | 106.62 |
| HEXACIMA | 104.14 |
| SYNFLORIX | 101.95 |
| TRITANRIX | 56.79 |
| ROTARIX | 36.35 |
| HEXYON | 36.08 |
| ROTATEQ | 31.32 |
| PREVENAR | 30.14 |
| M-M- RVAXPRO | 30.03 |
| NON | 29.97 |

Showing 1 to 10 of 41 entries

[Previous](#) [1](#) [2](#) [3](#) [4](#) [5](#) [Next](#)

So, who will we listen to, the **crying and screams of our own children**, or the propaganda pushed by pharma ?

VAERS on Sudden Infant Death

We can also look at VAERS data to see which vaccines have the strongest association with SIDS.

You can access the VAERS search tool here -

<https://knollfrank.github.io/HowBadIsMyBatch/SymptomsCausedByVaccines/index.html>

This tool covers data for 100 different vaccines, and the number shown is the lower confidence interval of the PRR. Enter the symptom “sudden infant death syndrome”

Select Symptom:

Show entries Search:

| Vaccine | Lower Confidence Limit of Proportional Reporting Ratio |
|---------|--|
| DTP | 65.3004187583126 |
| HEP | 42.58995655735134 |
| PNC | 22.268594913834622 |
| HIBV | 12.101787857542464 |
| MMR | 2.5465153291340243 |

Showing 1 to 5 of 5 entries Previous 1 Next

Select Symptom:

Show entries Search:

| Vaccine | Lower Confidence Limit of Proportional Reporting Ratio |
|------------|--|
| DTAPIPVHIB | 46.541323502336 |
| DTPHIB | 33.065371970797095 |
| DTP | 32.81742786754642 |
| DTAPHEPBIP | 24.127362813349265 |
| PNC | 16.83196236299044 |
| RV1 | 12.93736380445692 |
| HIBV | 8.037629376396943 |
| HEP | 6.839733165264602 |
| IPV | 5.750506613443228 |
| DTAP | 5.666145640192061 |

Showing 1 to 10 of 19 entries Previous 1 2 Next

Select Symptom:

Show entries

Search:

| Vaccine | Lower Confidence Limit of Proportional Reporting Ratio |
|------------|--|
| DTP | <u>37.98697813460069</u> |
| DTPHIB | <u>27.456127656928757</u> |
| PNC | <u>19.84058645735001</u> |
| 6VAX-F | <u>16.266383616898974</u> |
| MENB | <u>12.578727319724509</u> |
| DTAPIPVHIB | <u>11.04268074353193</u> |
| DTAP | <u>7.179043285839723</u> |
| IPV | <u>7.17630289801042</u> |
| RV1 | <u>5.77131547759157</u> |
| MMR | <u>5.288753131314622</u> |

Showing 1 to 10 of 28 entries

[Previous](#) [1](#) [2](#) [3](#) [Next](#)

Select Symptom:

Show entries

Search:

| Vaccine | Lower Confidence Limit of Proportional Reporting Ratio |
|------------|--|
| IPV | <u>118.27590371998468</u> |
| DTAPIPVHIB | <u>25.35532375420755</u> |
| MENB | <u>23.19719801062999</u> |
| RV1 | <u>9.30531954316926</u> |
| PNC | <u>7.525972707328947</u> |
| DTAP | <u>4.257004232872787</u> |
| MMR | <u>3.829266988605494</u> |
| PNC13 | <u>2.359847708887105</u> |

Showing 1 to 8 of 8 entries

[Previous](#) [1](#) [Next](#)

Select Symptom:

Show entries

Search:

| Vaccine | Lower Confidence Limit of Proportional Reporting Ratio |
|---------|--|
| PNC | 117.32827972532525 |
| PNC13 | 27.81215960902299 |

Showing 1 to 2 of 2 entries

Previous 1 Next

Vaccines and sudden infant death: An analysis of the VAERS database 1990-2019 and review of the medical literature

<https://pubmed.ncbi.nlm.nih.gov/34258234/>

“Although there is considerable evidence that a subset of infants has an increased risk of sudden death after receiving vaccines, health authorities eliminated "prophylactic vaccination" as an official cause of death, so medical examiners are **compelled to misclassify** and conceal vaccine-related fatalities under alternate cause-of-death classifications.

In this paper, the Vaccine Adverse Event Reporting System (VAERS) database was analyzed to ascertain the onset interval of infant deaths post-vaccination.

Of 2605 infant deaths reported to VAERS from 1990 through 2019, **58 % clustered within 3 days post-vaccination and 78.3 % occurred within 7 days post-vaccination, confirming that infant deaths tend to occur in temporal proximity to vaccine administration. The excess of deaths during these early post-vaccination periods was statistically significant ($p < 0.00001$).**

A review of the medical literature substantiates a link between vaccines and sudden unexplained infant deaths. Several theories regarding the pathogenic mechanism behind these fatal events have been proposed, including the role of inflammatory cytokines as neuromodulators in the infant medulla preceding an abnormal response to the accumulation of carbon dioxide; fatal disorganization of respiratory control induced by adjuvants that cross the blood-brain barrier; and biochemical or synergistic toxicity due to multiple vaccines administered concurrently. While the findings in this paper are not proof of an association between infant vaccines and infant deaths, they are highly suggestive of a causal relationship.

| Onset interval post-vaccination | Events reported | Cumulative % of total events |
|---------------------------------|-----------------|------------------------------|
| Day of Vaccination | 440 | 16.9 % (440/2605) |
| Day 2 | 760 | 46.1 % (1200/2605) |
| Day 3 | 312 | 58.0 % (1512/2605) |
| Day 4 | 214 | 66.3 % (1726/2605) |
| Day 5 | 131 | 71.3 % (1857/2605) |
| Day 6 | 92 | 74.8 % (1949/2605) |
| Day 7 | 92 | 78.3 % (2041/2605) |
| Days 8–60 | 564 | 100 % (2605/2605) |
| Total deaths | 2605 | |

Table shows how infant deaths decrease exponentially with each day after vaccination.

<https://pubmed.ncbi.nlm.nih.gov/34258234/>

NEWS

Most SIDS deaths occur within 48 hours of vaccination

“Sudden Infant Death Syndrome (SIDS) happens to babies, from 1 month to 1 year in age, but most frequently between 2 and 4 months of age.

A former police detective who handled over 250 SIDS cases, revealed that 50% of the SIDS cases happened within 48 hours after a vaccine was given and about 70% of the cases happened within one week of a vaccine.

This can only mean one thing: that the childhood vaccines are the leading cause of SIDS deaths.”

<https://vigilantnews.com/post/former-major-city-police-detective-reveals-50-of-sids-cases-happened-within-48-hours-post-vaccine>

Louisiana House Resolution

A House Resolution in Louisiana has been passed, urging the Department of Health to investigate the past 10 years of Sudden Infant Death Syndrome (SIDS) cases in the state and to analyze each report in relation to the infant's vaccination history.

TGA hides from questions about sudden infant deaths after vaccination

<https://blog.maryannedemasi.com/p/tga-hides-from-questions-about-sudden>

A string of sudden unexpected deaths in infants following the Infanrix Hexa® vaccine has forced the drug regulator to go to ground. Maryanne Demasi, PhD - Oct 28, 2024

Infanrix hexa and sudden death: a review of the periodic safety update reports submitted to the European Medicines Agency

https://www.researchgate.net/publication/319707034_Infanrix_hexa_and_sudden_death_a_review_of_the_periodic_safety_update_reports_submitted_to_the_European_Medicines_Agency

Periodic safety update reports (PSURs)

<https://www.ema.europa.eu/en/human-regulatory-overview/post-authorisation/pharmacovigilance-post-authorisation/periodic-safety-update-reports-psurs>

Vaccines are the real cause of Sudden Infant Death Syndrome

The term SIDS itself, according to @MidwesternDoc, is a euphemism used to cover up vaccine-related deaths.

One disturbing study found that two-thirds of infants who died of SIDS had been vaccinated with the DPT vaccine prior to death. Of those, 6.5% died within 12 hours of vaccination, 13% within 24 hours, and 26% within three days.

Dr. William Torch, the author of the study, concluded, "DPT vaccination may be a generally unrecognized major cause of sudden infant and early childhood death, and that the risks of immunization may outweigh its potential benefits."

@MidwesternDoc claims there is a century of evidence to suggest vaccines cause SIDS, but the government has continuously worked to suppress such findings.

One of those efforts has been the removal of vaccine-related death classifications in the ICD system in 1979. This change made it impossible for doctors to officially attribute infant deaths to vaccines, further obscuring the true cause of SIDS.

"The children who died from SIDS and their parents deserve recognition and justice," says @MidwesternDoc. "Infants cannot speak up for themselves. When you observe these vaccine injuries and the trauma they experience, it's very apparent what happened."

The Real Cause of Sudden Infant Death Syndrome Has Finally Been Revealed

The "conspiracy theorists" were right again.

<https://vigilantnews.com/post/the-real-cause-of-sudden-infant-death-syndrome-has-finally-been-revealed/>

Routine childhood vaccines cause sudden infant death syndrome (SIDS)

<https://expose-news.com/2024/07/12/vaccines-cause-sudden-infant-death-syndrome-sids/>

Twin Babies Die of 'Sudden Infant Death' 2 Days After Baby Shots

Twin girls, 3.5 months healthy babies with no medical history received 5 "baby wellness Shots"; Polio, Diphtheria, Tetanus, Pertussis, and Hepatitis B. Two days later their mother found them both dead in their crib.

483% INCREASE- SUDDEN INFANT DEATH SYNDROME (SIDS) - Nevada; 2019-22

Dr. James Thorp

- 2017 : 3.14
- 2018 : 0
- 2019 : 9.07
- 2020 : 12.11
- 2021 : 44.53
- 2022 : 52.91

<https://twitter.com/jathorpmfm/status/1727038583289037241?s=21&t=AO4hq7elhvP5MEcFEiaHRQ>

There is a century of evidence linking vaccines to sudden infant death

It was predicted that lockdowns would lead to a decline in SIDS due to reduced childhood vaccination and that's exactly what happened.

“A positive correlation between the number of vaccine doses and infant mortality rates (IMRs) is detectable in the most highly developed nations.”

<https://journals.sagepub.com/doi/10.1177/09603271111407644>

Read Steve Kirsch Substack (https://open.substack.com/pub/stevekirsch/p/more-vaccines-higher-infant-mortality?r=ytl4p&utm_medium=ios&utm_campaign=post)

11 Babies Died of SIDS within days

DTP vaccine was administered. <https://www.bitchute.com/video/Jy52tEnGCUIN>

NURSING 95 DRUG HANDBOOK

DPT or DTP vaccine side effects are listed in the Nursing 95 Drug Handbook ©1995 by Spring House Corporation.

VACCINATION ROULETTE – EXPERIENCES, RISKS AND ALTERNATIVES

DPT - FROM THE NURSING 95 DRUG HANDBOOK

ACTION: Mechanism: Promotes active immunity to diphtheria, tetanus and pertussis by inducing production of antitoxins and antibodies.

ONSET: After third dose;

DURATION: 4 to 6 years;

INDICATIONS AND DOSAGE:
Primary immunisation-Children 6 weeks to 6 years:
0.5 ml I.M. 2 months apart for three doses and a fourth dose 1 year later. Booster is 0.5 ml I.M. when starting school. Not advised for adults or children over 6 years. The acellular vaccine may be used only for the fourth or fifth dose in children 17 months to 7 years who have previously been immunized with three or four doses of the whole-cell vaccine.

ADVERSE REACTIONS

Systemic Common: anorexia, vomiting, slight fever, chills, malaise; Systemic, life-threatening: Seizures, encephalopathy, anaphylaxis, sudden infant death syndrome. Other: soreness, redness, expected nodule remaining several weeks at injection site.

Contraindications: Contraindicated in immuno-suppressed patients, in those on corticosteroid therapy, and in those with a history of seizures. Defer vaccination in patients with acute febrile illness. Children with pre-existing neurological disorders should not receive pertussis component. Also, children who exhibit neurologic signs after DPT injection shouldn't receive pertussis component in any succeeding injections. Diphtheria and tetanus toxoids should be given instead.

NURSING CONSIDERATIONS

Obtain history of allergies and reaction to immunisations. Acellular vaccine may be associated with a lower incidence of local pain and fever. DPT injection may be given at same time as trivalent oral polio vaccine.

Keep epinephrine [adrenalin] 1:1,000 available to treat anaphylaxis.

Do not use for active infection. Administer only by deep I.M. (intra-muscular) injection, preferably in the thigh or deltoid muscle. Don't give S.C. (subcutaneous, meaning under the skin). Shake before using. Refrigerate.

An information booklet is available that describes the risks and benefits of this vaccine. **MAKE SURE PARENTS READ AND UNDERSTAND THE INFORMATION BEFORE VACCINE IS ADMINISTERED.**

single antigen. In children under 6 years, use only when diphtheria, tetanus, and pertussis combination is contraindicated because of pertussis component.

NURSING CONSIDERATIONS

- Obtain history of allergies and reaction to immunization.
- Before injection, verify strength (pediatric or adult) of toxoid used.
- Keep epinephrine 1:1,000 available to treat anaphylaxis.
- Give in site not recently used for vaccines or toxoids.

diphtheria and tetanus toxoids and pertussis vaccine (DTP, DPT)

Tri-Immunol

diphtheria and tetanus toxoids and acellular pertussis vaccine

Acel-Imune

HOW SUPPLIED

whole-cell vaccine

Injection: 6.7 Lf (limit flocculation) units inactivated diphtheria, 5 Lf units inactivated tetanus, and 4 protective units pertussis per 0.5 ml, in 7.5-ml vials; 12.5 Lf units inactivated diphtheria, 5 Lf units inactivated tetanus, and 4 protective units pertussis per 0.5 ml, in 7.5-ml vials (Tri-Immunol)

acellular vaccine

Injection: 7.5 Lf units inactivated diphtheria, 5 Lf units inactivated tetanus, and 300 hemagglutinating units of acellular pertussis vaccine per 0.5 ml

ACTION

Mechanism: Promotes active immunity to diphtheria, tetanus, and pertussis by inducing production of anti-toxins and antibodies.

Onset: after third dose. **Duration:** 4 to 6 years.

INDICATIONS & DOSAGE

Primary immunization –

Children 6 weeks to 6 years: 0.5 ml I.M. 2 months apart for three doses and a fourth dose 1 year later. Booster is 0.5 ml I.M. when starting school.

Not advised for adults or children over 6 years.

The acellular vaccine may be used only for the fourth or fifth dose in children 17 months to 7 years who have previously been immunized with three or four doses of the whole-cell vaccine.

ADVERSE REACTIONS

Systemic: *seizures, encephalopathy, anorexia, vomiting, slight fever, chills, malaise, anaphylaxis, sudden infant death syndrome.*

Other: *soreness, redness, expected nodule remaining several weeks at injection site.*

INTERACTIONS

Immunosuppressive therapy: may reduce response to DPT vaccine. Avoid if possible.

CONTRAINDICATIONS

- Contraindicated in immunosuppressed patients, in those on corticosteroid therapy, and in those with a history of seizures. Defer vaccination in patients with acute febrile illness.
- Children with preexisting neurologic disorders should not receive pertussis component. Also, children who exhibit neurologic signs after DPT injection shouldn't receive pertussis component in any succeeding injections. Diphtheria and tetanus toxoids (DT) should be given instead.

NURSING CONSIDERATIONS

- Obtain history of allergies and reaction to immunization.
- Acellular vaccine may be associated

*Liquid form contains alcohol.

**May contain tartrazine.

Common reactions are in italics; *life-threatening*, in bold italics.

CLINICAL OBSERVATIONS

APNEA AS A WARNING SIGN OF SIDS

My safety signal analyses reveal that the same "vaccines" associated with SIDS are also associated with apnea and hypotonia. On further investigation I have found this finding confirmed by clinical literature.

"Respiratory observations made during nutritive feeding and a complete daytime nap within the first week of life on ten newborns who subsequently were victims of sudden infant death syndrome (SIDS) were compared with normative data based on 1,301 infants of comparable ages. Measurements were obtained of all apneic pauses and/or transient respiratory obstructive events at least two seconds in duration. A statistically significant increased number of future SIDS victims were found to have an unusual amount of apnea/obstruction episodes during nutritive feeding when compared with the normative group. Future SIDS victims, as a group, also had an increased frequency of apneic pauses during sleep. Furthermore, all SIDS victims demonstrated either an unusual amount of apnea/obstruction during nutritive feeding or apneic pauses during sleep. None of the airway obstruction measures during sleep differentiated the SIDS infants from the normative group. These results are consistent with the general hypothesis that prolonged apnea or airway obstruction is part of the pathophysiologic process resulting in SIDS and those hypotheses implicating unstable respiratory activity during sleep as well as apnea and pharyngeal/laryngeal dysfunction induced by liquid stimulation of the upper airway. They are also compatible with the growing body of evidence indicating that factors that predispose an infant to SIDS originate in the perinatal period. "

<https://pubmed.ncbi.nlm.nih.gov/7145538/>

What this means is that a simple breath monitor (which dings an alarm when a baby has an episode of apnoea greater than 20 seconds), will tell parents if their child is in danger of SIDS.

Apnoea is itself a manifestation of hypotonia of the respiratory system, and is also strongly associated with the suppression of feeding (eating and drinking). These are all tell-tale signs .

The worst response is to blame the parents . They are distraught enough with the death of a child, but to then claim they killed their child is so wrong that it borders on pure evil.

These findings establish preconditions of apnoea, hypotonia and diminished feeding that PRECEDE the fatal events of SIDS, thereby showing that the parents are NOT the cause. So, even in cases where SIDS inevitably occurs, all parents should monitor their "vaccinated" infant for apnoea and report every instance - because it will protect them

from accusations of homicide which might affect their parental status, and employment status later on.

I would also recommend taking a picture / video of any episodes of spasm or floppy infant - not least because it can be shown to a doctor or court later. The spasms can be so severe that they cause the body to arch so that only the heels and back of head are touching the floor, and the body contorts into a crescent moon shape. This may be accompanied by a high-pitched scream...

Close monitoring will also enable you to pinpoint the moment of onset of symptoms establishing a finer degree of temporal association with its cause (vaccine)

AUTISM AND SIDS

In addition to apnea, hypotonia, spasms and suppressed feeding, SIDS is also associated with autism. So these are the constellations of symptoms that precede SIDS.

The drugs with highest incidence rates for autism are the very same drugs that have highest incidence rates for SIDS, confirming that SIDS is the result of catastrophic vaccine induced brain damage.

So Sudden Infant Death is not so sudden.

Rather, it is preceded by these constellations which provide a means of

1. Detecting vulnerability to SIDS
2. Protecting parents from false allegations

On the next page I used the vaccine search engine to compare the vaccines associated with SIDS with the vaccines associated with AUTISM. It turns out they are the same list. The profound consequence of this is that SIDS doesn't just happen, but is preceded by the entire constellation of symptoms that we associate with autism.

This furnishes us with yet another means of detecting vulnerability to SIDS before it happens.

Here are the vaccines associated with SIDS – you can access this list by using the search engine here - <https://knollfrank.github.io/HowBadIsMyBatch/SymptomsCausedByVaccines/index.html?symptom=Sudden+infant+death+syndrome>

Select Symptom:

Show entries Search:

| Vaccine | Lower Confidence Limit of Proportional Reporting Ratio |
|---------|--|
| DTP | 65.3004187583126 |
| HEP | 42.58995655735134 |
| PNC | 22.268594913834622 |
| HIBV | 12.101787857542464 |
| MMR | 2.5465153291340243 |

Showing 1 to 5 of 5 entries Previous 1 Next

And here is the list of vaccines associated with autism –

Select Symptom:

Show entries Search:

| Vaccine | Lower Confidence Limit of Proportional Reporting Ratio |
|---------|--|
| MMR | 207.29225277842704 |
| PNC | 12.245920960399944 |
| DTP | 8.882537712012407 |
| HIBV | 7.220608765969972 |
| DTAP | 6.778551145911666 |
| HEP | 5.411393752290311 |

Showing 1 to 6 of 6 entries Previous 1 Next

There is considerable overlap, demonstrating that SIDS is preceded by the entire constellation of symptoms associated with autism.

HOW TO SPOT A DANGER OF SIDS

1. Excess crying, screaming
2. Loss of appetite, loss of thirst
3. Episodes of breath holding, apnoea
4. Floppy infant, hypotonia
5. Spasms, convulsions
6. Autistic symptoms – loss of eye contact
7. Bulging fontanelle
8. Temporal proximity (danger period) within 7 days of a "vaccination"
9. Relative danger posed by different “vaccines”
 - a. DTP based vaccines have the highest risk
 - b. Multi-valent vaccines have the highest risk

Since cessation of breathing is likely to be the fatal effect, then some way of monitoring breathing would be useful.

Breathing monitors are available - <https://www.lullabytrust.org.uk/wp-content/uploads/Movement-Monitors-Fact-Sheet-December-2021-update.docx.pdf>

THE DANGER PERIOD

Another thing to remember is that Miller (<https://pubmed.ncbi.nlm.nih.gov/34258234/>) found that SIDS events are clustered within the first 7 days after a vaccination. So you need to monitor your baby closely within this time - even if it means taking shifts. If your baby stops breathing you will need to know how to administer CPR.

- Day 1 17%
- Day 2 30%
- Day 3 12%
- Day 4 8 %
- Day 5 5%
- Day 6 3%
- Day 7 3%

78% of SIDS events occur within the first 7 days.

COMMITMENT TO MONITOR

Since “vaccination” puts a child at risk of SIDS, then a decision to “vaccinate” must include a parental commitment to monitor the child for at least 7 days after administration of the “vaccine”.

COMMITMENT TO CARE

Since “vaccination” also puts a child at risk of BRAIN DAMAGE, as seen in dysregulation of breathing, hunger and thirst reflexes, spasms, convulsions, hypotonia, and brain inflammation, then a decision to “vaccinate” must include a parental commitment to care for an autistic child for the remainder of that child’s life.